WPO36771

Sandia National Laboratories

Albuquerque, New Mexico 87185

date: 29 March 1996

to: Christine T. Stockman, MS-1328 (Org. 6749)

Hans W. Papenguth

from: Hans W. Papenguth, MS-1320 (Org. 6748)

subject: Colloidal Actinide Source Term Parameters

This memorandum summarizes best estimates for the mobile colloidal actinide source term for input to the WIPP Compliance Certification Application. The use of material and parameter identification codes is consistent with your letter to me dated 29 March 1996 requesting parameter values. In the attached table, I have provided best estimates for the following material-parameter combinations:

IDMTRL: Th, U, Np, Pu, Am

IDPRAM:

CONCINT	concentration of actinide associated with mobile actinide intrinsic colloids
CONCMIN	concentration of actinide associated with mobile mineral fragment colloids
CAPHUM	maximum concentration of actinide associated with mobile humic colloids
CAPMIC	maximum concentration of actinide associated with mobile microbes
PROPHUM	proportionality constant for concentration of actinides associated with mobile humic colloids
PROPMIC	proportionality constant for concentration of actinides associated with mobile microbes

As a first approximation, the colloidal behavior of curium can be simulated be using parameter values for americium. The basis for the values summarized in the attached table is described in the following record packages for WBS 1.1.10.2.1:

WPO#	Parameter Record Package Name
35850	Mobile-Colloidal-Actinide Source Term. 1. Mineral Fragment Colloids
35852	Mobile-Colloidal-Actinide Source Term. 2. Actinide Intrinsic Colloids
35855	Mobile-Colloidal-Actinide Source Term. 3. Humic Substances
35856	Mobile-Colloidal-Actinide Source Term. 4. Microbes

Information Only

copy to:

MS 1328	Hong-Nian Jow, 6741
MS 1328	Amy S. Johnson, 6741
MS 1328	Martin S. Tierney, 6741
MS 1320	E. James Nowak, 6831 EJN
MS 1320	R. Vann Bynum, 6831
MS 1341	John T. Holmes, 6748
MS 1341	Laurence H. Brush, 6748
MS 1341	Robert C. Moore, 6748
MS 1341	W. Graham Yelton, 6748
MS 1320	W. George Perkins, 6748 wgp
MS 1320	John W. Kelly, 6748
MS 1320	Daniel A. Lucero, 6748
MS 1320	Craig F. Novak, 6748
MS 1320	Hans W. Papenguth, 6748
MS 1320	Malcolm D. Siegel, 6748
MS 1324	Susan A. Howarth, 6115
MS 1341	Kurt O. Larson, 6747
MS 1341	Ruth F. Weiner, 6747
MS 1324	Richard Aguilar, 6851

SWCF-A:WBS1.1.10.2.1

Information Only

Parameter	Material	Most Likely Value	Minimum Value	Maximum Value	Units	Distribution Type	Notes
CONCMIN	Th	1.3e-09	1.3e-10	1.3e-08	moles colloidal mineral-fragment-bound Th per liter of dispersion	triangular	1
CONCMIN	U	1.3e-09	1.3e-10	1.3e-08	moles colloidal mineral-fragment-bound U per liter of dispersion	triangular	1
CONCMIN	Np	1.3e-09	1.3e-10	1.3e-08	moles colloidal mineral-fragment-bound Np per liter of dispersion	triangular	1
CONCMIN	Pu	1.3e-09	1.3e-10	1.3e-08	moles colloidal mineral-fragment-bound Pu per liter of dispersion	triangular	1
CONCMIN	Am	1.3e-09	1.3e-10	1.3e-08	moles colloidal mineral-fragment-bound Am per liter of dispersion	triangular	1
CONCINT	Th	0.0e+00	0.0e+00	0.0e+00	moles actinide-intrinsic colloidal Th per liter of dispersion	constant	
CONCINT	U	0.0e+00	0.0e+00	0.0e+00	moles actinide-intrinsic colloidal U per liter of dispersion	constant	
CONCINT	Np	0.0e+00	0.0e+00	0.0e+00	moles actinide-intrinsic colloidal Np per liter of dispersion	constant	
CONCINT	Pu	1.0e-09	1.0e-09	1.0e-09	moles actinide-intrinsic colloidal Pu per liter of dispersion	constant	
CONCINT	Am	0.0e+00	0.0e+00	0.0e+00	moles actinide-intrinsic colloidal Am per liter of dispersion	constant	
PROPHUM	Th	6.4e+00	6.4e+00	6.4e+00	moles colloidal humic-bound Th per moles dissolved Th	constant	2,3
PROPHUM	U	1.4e+00	1.6e-01	2.0e+00	moles colloidal humic-bound U per moles dissolved U	triangular	2,3,4
PROPHUM	Np	4.0e+00	4.0e+00	4.0e+00	moles colloidal humic-bound Np per moles dissolved Np	constant	2,3
PROPHUM	Pu	5.9e+00	5.9e+00	5.9e+00	moles colloidal humic-bound Pu per moles dissolved Pu	constant	2,3
PROPHUM	Am	2.5e+00	1.9e-01	3.9e+00	moles colloidal humic-bound Am per moles dissolved Am	triangular	2,3,4

Information Only

Parameter	Material	Most Likely Value	Minimum Value	Maximum Value	Units	Distribution Type	Notes
CAPHUM	Th	1.5e-05	1.5e-05	1.5e-05	moles colloidal humic-bound Th per liter of dispersion	constant	5,6
CAPHUM	U	1.5e-05	1.5e-05	1.5e-05	moles colloidal humic-bound U per liter of dispersion	constant	5,6
CAPHUM	Np	1.5e-05	1.5e-05	1.5e-05	moles colloidal humic-bound Np per liter of dispersion	constant	5,6
CAPHUM	Pu	1.5e-05	1.5e-05	1.5e-05	moles colloidal humic-bound Pu per liter of dispersion	constant	5,6
CAPHUM	Am	1.5e-05	1.5e-05	1.5e-05	moles colloidal humic-bound Am per liter of dispersion	constant	5,6
PROPMIC	Th	3.1e+00	3.1e+00	3.1e+00	moles microbial Th per moles dissolved Th	constant	2,3
PROPMIC	U	2.1e-03	2.1e-03	2.1e-03	moles microbial U per moles dissolved U	constant	2,3
PROPMIC	Np	1.2e+01	1.2e+01	1.2e+01	moles microbial Np per moles dissolved Np	constant	2,3
PROPMIC	Pu	3.0e-01	3.0e-01	3.0e-01	moles microbial Pu per moles dissolved Pu	constant	2,3
PROPMIC	Am	3.6e+00	3.6e+00	3.6e+00	moles microbial Am per moles dissolved Am	constant	2,3
CAPMIC	Th	1.9e-03	1.9e-03	1.9e-03	moles total mobile Th per liter	constant	5,7
CAPMIC	U	2.1e-03	2.1e-03	2.1e-03	moles total mobile U per liter	constant	5,7
CAPMIC	Np	2.7e-03	2.7e-03	2.7e-03	moles total mobile Np per liter	constant	5,7
CAPMIC	Pu	6.8e-05	6.8e-05	6.8e-05	moles total mobile Pu per liter	constant	5,7
CAPMIC	Am	not currently available	not currently available	not currently available	moles total mobile Am per liter	constant	5,7

Notes:

general	The colloidal actinide source term is added to the dissolved actinide source term.
general	None of the parameters are correlated.
1	If a distribution is not used for mineral-fragment-bound actinides, use the maximum concentration as a constant value.
2	Proportionality constants may be used with actinide solubility expressed in molarity or molality, depending on the desired final units.
3	Proportionality constants are to be used with the sum of actinide oxidation species for each actinide element (uncomplexed only, i.e., without organic ligand contribution).
4	If a distribution is not used for humic-bound U or Am, use the maximum concentration as a constant value.
5	The maximum ("cap") values are in units comparable to molarity rather than molality.
6	CAPHUM is compared to the concentration of the respective humic-bound actinide element.
7	CAPMIC is compared to the total concentration of the respective actinide element in the mobile system (i.e. the sum of dissolved plus colloidal actinide).

Information Only